

Comments regarding
Dietary Guidelines for Americans

Submitted to the
Dietary Guidelines Advisory Committee,
U.S. Department of Health and Human Services, and
U.S. Department of Agriculture

January 23, 2009

Submitted to:
Crystal Tyler
Graduate School, USDA
600 Maryland Avenue, SW
Suite 330
Washington, D.C. 20024

The Center for Science in the Public Interest (CSPI) respectfully submits to the Dietary Guidelines Advisory Committee (DGAC), the U.S. Department of Health and Human Services (DHHS), and the U.S. Department of Agriculture (USDA) recommendations regarding the bulletin *Nutrition and Health: Dietary Guidelines for Americans*.

CSPI is a non-profit consumer education and advocacy organization that since 1971 has been working to improve the public's health through better nutrition and safer food. CSPI's work is supported primarily by its 800,000 members and subscribers to its *Nutrition Action Healthletter*, the nation's largest circulation health newsletter. CSPI does not accept any government or corporate funding.

CSPI's work was instrumental in passage of the Nutrition Labeling and Education Act of 1990 and the Alcoholic Beverage Labeling Act of 1988. Other initiatives include studies of the nutritional quality of restaurant foods, advocating trans fat labeling on packaged foods, and campaigns to promote low-fat milk consumption, improve school foods, stop misleading food and alcohol advertising, enforce food safety laws, and improve alcoholic-beverage labeling.

Enclosed are eight sets of comments regarding the following guidelines:

- Nutrient Adequacy
- Sodium
- Fibers
- Whole Grains
- Added Sugars
- Energy Balance
- Fatty Acids
- Restaurant Foods
- Food Dyes and Behavior
- Ethanol

Our comments are summarized in the oral testimony which will be presented to the committee on January 29, 2009, which is enclosed.

For more information or questions regarding these comments please contact Alexandra Lewin, Ph.D. at 202.777.8351 or alewin@cspinet.org.

Comments by the Center for Science in the Public Interest on Sodium

Salt – sodium chloride – is perhaps the deadliest ingredient in our food supply.

The health community is united in the conclusion that Americans are consuming far too much salt and other sources of sodium. Consuming less sodium is one of the single most important ways to prevent cardiovascular disease. The salt guideline, while currently quite good, should do even more to alert Americans to: the risks of consuming excess amounts of sodium, how much sodium is included in processed and restaurant foods, and the daily limit of sodium.

Sixty-five million Americans have high blood pressure,¹² another 45 million people have “prehypertension,”¹³ and about 90 percent of Americans will eventually develop hypertension.¹⁴ African Americans’ rate of hypertension is 60 percent greater, and rate of stroke deaths is 40 percent greater, than that of the general population.¹⁵

In contrast to the U.S. Government, the United Kingdom’s Food Standards Agency (FSA) has made salt reduction a top priority and is both making consumers more aware of the health threat posed by excessive salt consumption and exerting strong pressure on the food and restaurant industries to gradually lower sodium levels to specified targets. The DGAC should provide an authoritative statement that would encourage our government, too, to make salt reduction a top health priority.

Despite experts’ admonitions over the years, per capita sodium consumption has actually increased (according to NHANES surveys) from 2,800 mg in 1976-80 to 3,400 mg in 2003-04. According to the 2003-04 NHANES survey, 70 percent of adult males and 50 percent of adult females exceed the 2,300 mg Dietary Guidelines recommended level—even though those estimates omit salt added at the table or in cooking, and the NHANES respondents are thought to underestimate their actual consumption. Thus, actual daily consumption is probably closer to 4,000 mg,¹⁶ with high percentages of both males and females exceeding the recommended levels.

The extraordinary importance of lowering sodium consumption was highlighted in a 2004 commentary in the *American Journal of Public Health* coauthored by Claude Lenfant, then-Director of NHLBI, and two colleagues. The article estimated that reducing the sodium content of packaged and restaurant foods by 50 percent would prevent 150,000 deaths due cardiovascular disease per year.¹⁷ Reducing sodium consumption also would save tens of billions of dollars in health care costs. A preliminary RAND Corp. study estimates that reducing average sodium from 3,400 mg to 2,300 mg per day would reduce direct medical costs by \$18 billion per year and quality of life by \$32 billion per year.¹⁸ A further reduction to 1,500 mg per day would reduce medical costs by \$28 billion per year.

In the early 1980s, the FDA called for vigorous voluntary action on the part of industry to reduce sodium levels. We evaluated the changes in almost 100 foods that were produced in 1983 and still produced in 2004.¹⁹ The average change in sodium content of those foods was only a 5 percent decrease, a decline of 0.3 percent per year—a far lower rate than what would be needed to achieve the 50-percent decrease over 10 years that the American Medical Association, American Heart Association, American Public Health Association, and many other experts have recommended.²⁰

In 2005, CSPI evaluated the sodium content of 528 randomly chosen manufactured and restaurant foods. Notwithstanding the 2005 Dietary Guidelines' strong admonition to consume less salt, when we surveyed the same foods in late 2008, we found that the average sodium content changed (increased) by less than one percent. Clearly, the food industry needs to be sent a stronger signal that major sodium reductions are essential.

I. The guidelines should explicitly state the direct link of sodium consumption to heart attacks and death

Not only is sodium in the American diet a major cause of high blood pressure, but there is now hard evidence that curbing sodium also reduces the risk of cardiovascular disease (heart attack, stroke, coronary bypass, angioplasty, or cardiovascular death). According to a recent study of more than 3,000 overweight people aged 30 to 54 with pre-hypertension, those who were randomly assigned to reduce their sodium intake (the average reduction was roughly 800 to 1,000 milligrams a day) for 1½ to 3 years were 30 percent less likely to be diagnosed with cardiovascular disease over the next 10 to 15 years.²¹ Also, they reduced their risk of dying from cardiovascular disease by 20 percent, although that data was not statistically significant. Furthermore, when researchers gave salt tablets (4,600 mg of sodium) to 16 young healthy normotensives who were consuming a low-sodium diet (600 mg a day) for five days, vascular endothelial function, left ventricular mechanical relaxation, and electric repolarization were impaired.²² Also, in a study of elderly Taiwanese military veterans living in a retirement home, researchers replaced regular salt with potassium-enriched salt, thereby boosting the men's potassium intake while lowering their sodium intake.²³ After almost three years of follow-up, those men who reduced their sodium intake enjoyed a 41-percent reduction in cardiovascular disease deaths. The guidelines should explicitly state that there is a direct link not only between sodium consumption and blood pressure, but also between sodium and cardiovascular disease.

II. The guidelines should lower the recommendation of daily consumption of sodium for most adults to 1,500 mg.

The 2005 Dietary Guidelines recommends that people limit consumption of sodium to 2,300 mg per day unless they are in certain population groups. Individuals with hypertension, blacks, and middle-aged and older adults are advised to limit their sodium to 1,500 mg per day. Because about two-thirds of the population is in one or another of those “specific population groups,”²⁴ the recommendation of daily consumption of

sodium by adults should be lowered to 1,500 mg. Lower limits should be indicated for certain other population groups, such as 1,300 mg for people 51 to 70 years old and 1,200 mg for those over 70. Those numbers reflect the Institute of Medicine's (IOM) 2004 Dietary Reference Intakes for daily "Adequate Intake" for sodium. The IOM offers two sets of recommendations, an adequate intake and an upper level intake. But according to the IOM, "...the UL is not a recommended intake and, as with other ULs, there is no benefit to consuming levels above the AI."²⁵

III. The Guidelines should include a specific sodium recommendation for children.

The 2005 Sodium and Potassium guideline recommends intakes of potassium, but not sodium, for children. (Currently, recommendations for sodium intake by children are only included in the 2005 DGAC report – not the Dietary Guidelines itself - and are higher than those recommended by the IOM.) The IOM's recommended adequate intake of sodium is 1,000 mg/day for children aged 1 to 3, 1,200 mg/day for children aged 4 to 8, and 1,500 mg/day for children aged 9 to 13.²⁶ Children, however, are consuming an average of more than twice the recommended amounts of sodium.²⁷

A high-salt diet in childhood is associated with higher blood pressure in children that likely will lead to increased risks of hypertension, strokes and heart disease later in life.²⁸ Reducing the sodium content in kids' foods would help train their taste buds to enjoy lower-salt foods. In addition, some researchers have attributed the rise in the rate of kidney stones seen in children to the excess amounts of salt in children's diets.

CSPI recently conducted a study of the nutritional quality of 1,474 kids' meals at 13 top chain restaurants and found that not only are those meals loaded with calories, but that every single meal was too high in sodium (based on 1/3 of the IOM's recommended adequate intake for children).²⁹ In fact, many of the meals included over a whole day's worth of sodium. For example, the KFC kids' meal that includes crispy chicken strips, green beans, baked beans, Tropicana fruit punch, and Teddy Grahams supplies 2,215 mg of sodium. At Denny's, the Galactic Grilled Cheese and Moon Crater Mashed Potatoes with Gravy combo meal has 1,385 mg of sodium. Chili's offers a grilled chicken sandwich, rice, and 1 % chocolate milk meal that contains 1,610 mg of sodium. At McDonald's, a child can order a meal that includes a double cheeseburger, small French fries, and a 12 oz. fountain drink meal - and 1,340 mg of sodium.

The guidelines should make specific recommendations for children's intake of sodium and also highlight the need to pay careful attention to the amounts of sodium consumed by children outside of the home.

IV. The sodium guideline should emphasize the extraordinarily high levels of sodium in many restaurants foods.

Although the 2005 guideline does state that "...food served by food establishments may be high in sodium," it does not emphasize sufficiently the extraordinarily high levels of

sodium in many restaurant foods. The sodium content of foods listed in Table 15 of the Guidelines does not include a single restaurant meal. The Dietary Guidelines should provide more details regarding high-sodium foods in restaurants, especially given that one-third of all calories are consumed away from home. According to CSPI's lab analyses:³⁰

- Typical deli sandwiches have 1,000 mg or more of sodium. Some, like corned beef with mustard, ham with mustard or mayo, or a turkey club, contain closer to 2,000 mg of sodium. A Reuben or “overstuffed corn beef” sandwich can contain closer to 3,000 mg of sodium.
- Nearly all main courses at popular Italian restaurants had at least 1,000 to 2,000 mg of sodium. Most Chinese dishes and Mexican platters had at least 2,000 to 3,000 mg of sodium.
- Many fast foods include high amounts of sodium. Breakfast sandwiches include 1,000 mg of sodium or more. Also, many grilled chicken sandwiches and large cheeseburgers have well over 1,000 mg of sodium.
- Many appetizers at popular dinner house chains are loaded with sodium. For example, an order of buffalo wings or fried mozzarella sticks has approximately 2,000 mg of sodium.

If the Dietary Guidelines for Americans is going to provide useful information to consumers, it is essential that it give practical advice for eating in restaurants and other food service establishments where Americans get about one-third of their calories.

V. Recommended research.

We urge the committee to recommend that new research be sponsored to better estimate sodium consumption and to monitor changes in consumption.

- It is important to know how much sodium is consumed from packaged foods, restaurant foods, salt added during cooking, and salt added at the table. A widely cited report was conducted decades ago involving only 62 adults who were not nationally representative.³¹ The committee should recommend that the FDA, CDC, or NHLBI conduct a new study to more accurately identify the sources of sodium in the American diet.
- Health officials (and the DGAC) do not have access to reliable data on sodium consumption. Purported per capita sodium consumption is deduced from dietary recalls conducted by NHANES. Unfortunately, the NHANES methodology includes significant flaws: food intake is typically underestimated, brand-to-brand variations (in both restaurant and packaged foods) are large, and salt added to foods by consumers is not included. Hence, the widely used NHANES figures probably underestimate actual consumption by 15 percent or more.³² Urinary-

excretion studies are the only accurate means of determining total sodium consumption, but, unfortunately, only small, non-representative studies are available. The DGAC should recommend that 24-hour urinary-excretion measurements be added to the NHANES study protocols.

-
- 1 Appel LJ, et. al. Effects of protein, monounsaturated fat, and carbohydrate intake on blood pressure and serum lipids: results of the OmniHeart randomized trial. *JAMA*. 2005 Nov 16;294(19):2455-64.
- 2 Jacobson, MF. Six Arguments for a Greener Diet. Washington, DC (Center for Science in the Public Interest). 2006.
- 3 21 CFR 101.14 (e) (6)
- 4 Nutrition Action Healthletter June 2008, page 1.
- 5 *Cancer*. 2008 Jul 1;113(1):150-7.
- 6 *Cochrane Database Syst Rev*. 2008 Apr 16;(2):CD007176.
- 7 *Nutr Clin Pract*. 2008 Oct-Nov;23(5):468-76.
- 8 *J Natl Cancer Inst*. 2009 Jan 7;101(1):14-23.; *JAMA*. 2008 Nov 12;300(18):2123-33.
- 9 *Cochrane Database Syst Rev*. 2008 Apr 16;(2):CD007176.
- 10 *Curr Atheroscler Rep*. 2008 Dec;10(6):503-9
- 11 *Heart*. 2006 Feb;92(2):166-9.
- 12 Fields LE, Burt VL, Cutler JA, et al. The burden of adult hypertension in the United States 1999 to 2000: a rising tide. *Hypertension*. 2004; 44:398-404.
- 13 Chobanian AV, Bakris GL, Black HR, et al. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *JAMA*. 2003; 289:2560-72. National Heart, Lung, and Blood Institute. NHLBI issues new high blood pressure clinical practice guidelines. News release. May 14, 2003.
- 14 Vasan RS, Beiser A, Shisadri S. Residual lifetime risk for developing hypertension in middle-aged women and men. *JAMA*. 2002; 287:1003-10.
- 15 National Center for Health Statistics. *Health, United states, 2004, With Chartbook on Trends in the Health of Americans*. Hyattsville, MD, 2004.
- 16 Zhou BF, Stamler J, Dennis B, et al. Nutrient intakes of middle-aged men and women in China, Japan, United Kingdom, and United States in the late 1990s: The INTERMAP study. *J Hum Hypertens*. 2003; 17:623–30. Kumanyika SK, Cook NR, Cutler JA, et al. Sodium reduction for hypertension prevention in overweight adults: further results from the Trials of Hypertension Prevention Phase II. *J Hum Hypertens*. 2005; 19:33–45.
- 17 Havas, S, Rocella EJ, Lenfant C. Reducing the public health burden from elevated blood pressure levels in the United States by lowering intake of dietary sodium. *Am J Pub Health*. 2004; 94:19-22. Other studies, when extrapolated to the United States, also indicate that halving sodium would save 100,000 to 200,000 lives per year: He FJ, MacGregor GA. How far should salt intake be reduced? *Hypertension*. 2003; 42:1093–9. (CSPI averaged the percentage reductions based on systolic and diastolic blood pressures.)

Tuomilehto J, Jousilahti P, Rastenyte D, et al. Urinary sodium excretion and cardiovascular mortality in Finland: a prospective study. *Lancet*. 2001;357:848–51.

18 Palar, K, Sturm, R. The Benefits of Reducing Sodium Consumption in the US Adult Population. Rand Health, Academy Health Annual Research Meeting Presentation, June 9, 2008.

19 Jacobson MF. Sodium content of processed foods: 1983–2004. *Am J Clin Nutr*. 2005;81:941a-2a.

20 Chobanian AV, Bakris GL, Black HR, et al. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *JAMA*. 2003; 289:2560-72.

21 Cook NR, Cutler JA, Obarzanek E, et al. Long-term effects of dietary sodium reduction on cardiovascular disease outcomes: Observational follow-up of the trials of hypertension prevention (TOHP). *BMJ*. 2007; 334: 885-93.

22 Tzemos N, et al. Adverse cardiovascular effects of acute salt loading in young normotensive individuals. *Hypertension*. 2008 Jun;51(6):1525-30.

23 Chang HY, Hu YW, Yue CS, et al. Effect of potassium-enriched salt on cardiovascular mortality and medical expenses of elderly men. *Am J Clin Nutr*. 2006 Jun;83(6):1289-96.

24 Labarthe, D. (2009, January). Presentation at the IOM Meeting on Strategies to Reduce Sodium Intake, Washington, D.C.

25 Institute of Medicine: “Dietary Reference Intakes for Water, Potassium, Sodium, Chloride, and Sulfate.” Washington DC: National Academy Press, p 271, 2004.

26 Institute of Medicine: “Dietary Reference Intakes for Water, Potassium, Sodium, Chloride, and Sulfate.” Washington DC: National Academy Press, p 307, 2004.

27 NHANES, 2001-02. “What We Eat in America.” <http://www.ars.usda.gov/foodsurvey>

28 Feng J. He; Graham A. MacGregor. Importance of Salt in Determining Blood Pressure in Children. *Hypertension*. 2006; 48:861.

29 Wootan M, Batada A, Marchlewicz E. Kids Meals: Obesity on the Menu. Center for Science in the Public Interest: Washington, D.C.; 2008.

30 Jacobson MF, Hurley J. Restaurant Confidential. New York: Workman, 2002.

31 Mattes RD, Donnelly D. Relative contributions of dietary sodium sources. *J Am Coll Nutr*. 1991;10:383–93.

32 The actual level is probably close to 4,000 mg. Zhou BF, Stamler J, Dennis B, et al. Nutrient intakes of middle-aged men and women in China, Japan, United Kingdom, and United States in the late 1990s: The INTERMAP study. *J Hum Hypertens*. 2003; 17:623–30. Kumanyika SK, Cook NR, Cutler JA, et al. Sodium reduction for hypertension prevention in overweight adults: further results from the Trials of Hypertension Prevention Phase II. *J Hum Hypertens*. 2005;19:33–45.

33 Grabske HA, Slavin JL. Low-digestible carbohydrates in practice. *J Am Diet Assoc*. 2008 Oct;108(10):1677-81.

34 Food and Nutrition Board. Institute of Medicine. Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein and Amino Acids (National Academies Press, Washington, DC) 2005. p. 362.

-
- 35 Elaine Wong, "Fiber Being Pitched As Tasty, Sexy Even." *Brandweek*. Jan. 17, 2009.
- 36 Home and Garden Bulletin No. 252. USDA. Aug. 1992, slightly revised Oct. 1996. p. 17.
- 37 WHO-FAO. Diet, Nutrition and the Prevention of Chronic Diseases. WHO Technical Report Series 916. 2003 (www.fao.org/docrep/005/AC911E/AC911E00.HTM (accessed Jan. 10, 2009)).
- 38 Dietary Guidelines for Americans. www.health.gov/dietaryguidelines/dga2005/document/html/appendixA.htm (accessed Jan. 19, 2009). Appendix A-3.
- 39 Letter to Secretary Tommy Thompson, April 15, 2003.
- 40 Melanson KJ, Angelopoulos TJ, Nguyen V, et al. High-fructose corn syrup, energy intake, and appetite regulation. *Am J Clin Nutr*. 2008;88(suppl):1738S-44S.
- 41 Bray GA. How bad is fructose? *Am J. Clin Nutr*. 2007;86:895-6.
- 42 Havel PJ. Dietary fructose: implications for dysregulation of energy homeostasis and lipid/carbohydrate metabolism. *Nutr Rev*. 2005;63:133-57.
- 43 Stanhope KL, Havel PJ. Endocrine and metabolic effects of consuming beverages sweetened with fructose, glucose, sucrose, or high-fructose corn syrup. *Am J Clin Nutr*. 2008 Dec;88(6):1733S-7S.
- 44 ERS production data for HFCS-42 and HFCS-55, which the Corn Refiners Association says account for virtually all HFCS and crystalline fructose, indicate that the weighted-average sugar content of HFCS is almost exactly 50% glucose and 50% fructose. (HFCS-42: 3,660 (1,000 short tons, dry weight); HFCS-55: 5479 (1,000 short tons, dry weight))
- 45 Some products are sweetened with pure fructose (and sometimes bear labels claiming "No HFCS").
- 46 CSPI petition to FDA. July 13, 2005. www.cspinet.org/new/pdf/final_soda_petition.pdf.
- 47 Analyses by Environ. Calculations conducted in September 1998 for the Center for Science in the Public Interest by Environ, Inc., Arlington, Va., based on USDA CSFII 1994-96 Data Tables.
- 48 Giammattei J, Blix G, Marshak HH, et al. Television watching and soft drink consumption. *Arch Pediatr Adolesc Med*. 2003;157:882-6.
- 49 Troiano RP, Briefel RR, Carroll MD, Bialostosky K. Energy and fat intakes of children and adolescents in the United States: data from the National Health and Nutrition Examination Surveys. *Am J Clin Nutr*. 2000;72(suppl):1343S-53S.
- 50 Ludwig DS, Peterson KE, Gortmaker SL. Relationship between consumption of sugar-sweetened drinks and childhood obesity: a prospective, observational analysis. *Lancet*. 2001;357:505-8.
- 51 Berkey CS, Rockett HR, Field AE, et al. Sugar-added beverages and adolescent weight change. *Obes Res*. 2004;12:778-88.
- 52 Some of the same researchers subsequently found that weight gain in the same cohort of children was also associated with milk consumption. Berkey CS, Rockett HRH, Willett WC, et al. Milk, dairy fat, dietary calcium, and weight gain. *Arch Pediatr Adolesc Med*. 2005;159:543-550.
- 53 Schulze MB, Manson JE, Ludwig DS, et al. Sugar-sweetened beverages, weight gain, and incidence of type 2 diabetes in young and middle-aged women. *JAMA*. 2004;292:927-34.

54 Apovian CM. Sugar-sweetened soft drinks, obesity, and type 2 diabetes. *JAMA*. 2004;292:978-9.

55 Palmer JR, Boggs DA, Boggs MS, et al. Sugar-sweetened beverages and incidence of type 2 diabetes mellitus in African American women. *Arch Intern Med*. 2008;168:1487-92.

56 James J, Thomas P, Cavan D, et al. Preventing childhood obesity by reducing consumption of carbonated drinks: cluster randomized controlled trial. *BMJ*. 2004;328:1237.

57 Raben A, Vasilaras TH, Møller AC, et al. Sucrose compared with artificial sweeteners: different effects on ad libitum food intake and body weight after 10 wk of supplementation in overweight subjects. *Am J Clin Nutr*. 2002;76:721-9. An earlier, shorter study that compared soft drinks containing aspartame and high-fructose corn syrup found similar results. Tordoff MG, Alleva AM. Effect of drinking soda sweetened with aspartame or high-fructose corn syrup on food intake and body weight. *Am J Clin Nutr*. 1990;51:963-9.

58 Ebbeling CB, Feldman HA, Osganian SK, et al. Effects of decreasing sugar-sweetened beverage consumption on body weight in adolescents: a randomized, controlled pilot study. *Pediatrics*. 2006;117:673-80.

59 Malik VS, Schulze MB, Hu FB. Intake of sugar-sweetened beverages and weight gain: a systematic review. *Am J Clin Nutr*. 2006 Aug;84(2):274-88.

60 Vartanian LR, Schwartz MB, Brownell KD. Effects of soft drink consumption on nutrition and health: a systematic review and meta-analysis. *Am J Pub Health*. 2007;97(4):667-75.

61 Striegel-Moore RH, Thompson D, Affenito SG, et al. Correlates of beverage intake in adolescent girls: the National Heart, Lung, and Blood Institute Growth and Health Study. *J Pediatr*. 2006;148:183-7.

62 Dietz WH. Sugar-sweetened beverages, milk intake, and obesity in children and adolescents. *J Pediatr*. 2006;148:185-7.

63 Van Dam RM, Seidell JC. Carbohydrate intake and obesity. *Eur J Clin Nutr*. 2007;61 (suppl 1):S75-S99.

64 Mann J, Cummings JH, Englyst HN, et al. FAO/WHO scientific update on carbohydrates in human nutrition: conclusions. *Eur J Clin Nutr*. 2007;61 (suppl 1):S132-S7.

65 Committee on Prevention of Obesity in Children and Youth, Institute of Medicine. Fact Sheet, Parents can play a role in preventing childhood obesity. Sept. 2004.

66 DiMeglio DP, Mattes RD. Liquid versus solid carbohydrate: effects on food intake and body weight. *Intern J. Obesity*. 2000;24:794-800.

67 DeCastro JM. The effects of spontaneous ingestion of particular food or beverages on the meal pattern and overall nutrient intake of humans. *Physiol Behav*. 1993;53:1133-44. Mattes RD. Dietary compensation by humans for supplemental energy provided as ethanol or carbohydrate in fluids. *Physiol Behav*. 1996;59:179-87. Rolls BJ, Kim S, Federoff IC. Effects of drinks with sucrose or aspartame on hunger, thirst and food intake in men. *Physiol Behav*. 1990;48:19-26.

68 2005 Dietary Guidelines Advisory Committee Report, Part D, Section 2, pp. 18-9.

69 Almiron-Roig E, Drewnowski A. Hunger, thirst, and energy intakes following consumption of caloric beverages. *Physiol Behav*.

2003;79:767-73. DellaValle DM, Roe LS, Rolls BJ. Does the consumption of caloric and non-caloric beverages with a meal affect energy intake? *Appetite*. 2005;44:187-93. Such clinical studies cannot evaluate whether the drinking of particular beverages (because of their cultural uses, sizes, or tastes) changes what solid foods a person eats.

70 Ventura KA, Loken E, Birch LL. Risk profiles for metabolic syndrome in a nonclinical sample of adolescent girls. *Pediatrics*. 2006;118:2434-42.

71 Yoo S, Nicklas T, Baranowski T, et al. Comparison of dietary intakes associated with metabolic syndrome risk factors in young adults: the Bogalusa Heart Study. *Am J Clin Nutr*. 2004;80:841-8.

72 Dhingra R, Sullivan L, Jacques PF, et al. Soft drink consumption and risk of developing cardiometabolic risk factors and the metabolic syndrome in middle-aged adults in the community. *Circulation*. 2007;116:480-8.

73 Palmer JR, Boggs DA, Boggs MS, et al. Sugar-sweetened beverages and incidence of type 2 diabetes mellitus in African American women. *Arch Intern Med*. 2008;168:1487-92.

74 Docket No. FDA-2008-N-0040

75 Tucker KL, Morita K, Qiao N, et al. Colas, but not other carbonated beverages, are associated with low bone mineral density in older women: The Framingham Osteoporosis Study. *Am J Clin Nutr*. 2006;84: 936-42. Whiting SJ, Healey A, Psiuk S. Relationship between carbonated and other low nutrient dense beverages and bone mineral content of adolescents. *Nutr Res*. 2001;21:1107-15. Wyshak, G. Teenaged girls, carbonated beverage consumption, and bone fractures. *Arch Pediatr Adolesc Med*. 2000;154:610-3.

76 Saldana TM, Basso O, Darden R, et al. Carbonated beverages and chronic kidney disease. *Epidemiology*. 2007;18:501-6.

77 Choi HK, Curhan G. Soft drinks, fructose consumption, and the risk of gout in men: prospective cohort study. *BMJ* published online Jan 31, 2008;doi:10.1136/bmj.39449.819271.BE [accessed Feb. 4, 2008].

78 Choi JW, Ford ES, Gao X, et al. Sugar-sweetened soft drinks, diet soft drinks, and serum uric acid level: the Third National Health and Nutrition Examination Survey. *Arthritis Rheum*. 2008;59(1):109-16.

79 Rolls BJ, et al. Changing the energy density of the diet as a strategy for weight management. *J Am Diet Assoc*. 2005 May;105(5 Suppl 1):S98-103.

80 Ello-Martin JA, et al. Dietary energy density in the treatment of obesity: a year-long trial comparing 2 weight-loss diets. *Am J Clin Nutr*. 2007 Jun;85(6):1465-77

81 Savage JS, et al. Dietary energy density predicts women's weight change over 6 y.

Am J Clin Nutr. 2008 Sep;88(3):677-84.

82<http://mypyramid.gov/STEPS/nutrientdensefoods.html>.

83 Institute of Medicine. Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids. Washington, D.C.: National Academies Press, 2002.

84 American Heart Association. Heart and Stroke Statistics –2008 Update. Dallas, TX: American Heart Association, 2008. Accessed at

<http://www.americanheart.org/downloadable/heart/1200078608862HS_Stats%202008.final.pdf>

85 Clarke R. "Dietary lipids and blood cholesterol: quantitative meta-analysis of metabolic ward studies." *British Medical Journal* 1997, vol. 314, pp. 112-117.

-
- 86 Howell WH, et. al. "Plasma lipid and lipoprotein responses to dietary fat and cholesterol: a meta-analysis." *American Journal of Clinical Nutrition* 1997, vol. 65, pp. 1747-1764.
- 87 Armstrong ML, et. al. "Intimal thickening in normocholesterolemic rhesus monkeys fed low supplements of dietary cholesterol." *Circulation Research* 1974, vol. 34, pp. 447-454.
- 88 Walter Willett and Dariush Mozaffarian Ruminant or industrial sources of trans fatty acids: public health issue or food label skirmish? *Am. J. Clinical Nutrition*, Mar 2008; 87: 515 - 516.
- 89 CARDIOVASCULAR DISEASE RISK: Annie Motard-Bélanger, Amélie Charest, Geneviève Grenier, Paul Paquin, Yvan Chouinard, Simone Lemieux, Patrick Couture, and Benoît Lamarche. Study of the effect of trans fatty acids from ruminants on blood lipids and other risk factors for cardiovascular disease *Am. J. Clinical Nutrition*, Mar 2008; 87: 593 - 599.
- 90 Chardigny, Jet al Do trans fatty acids from industrially produced sources and from natural sources have the same effect on cardiovascular disease risk factors in healthy subjects? Results of the trans Fatty Acids Collaboration (TRANSFACT) study *Am. J. Clinical Nutrition*, Mar 2008; 87: 558 - 566.
- 91 The FDA permits such foods to contain less than 0.5 grams of trans fat per serving.
- 92 Some foods bragging "0g trans fat" contain substantial amounts of saturated fat.
93. 9 C.F.R. § 317.362 (e)(1)(2) (1998).
94. 59 Federal Register 26,916-26,917 (1994).
95. 9 C.F.R. § 317.362 (b)(6)(iv) (1998).
96. American Meat Institute. 2002 per capita disappearance data, personal communication, March 2004.
- 97 Lin B, Guthrie J, Frazao E. *Away-From-Home Foods Increasingly Important to Quality of American Diet*. Washington, DC: U.S. Department of Agriculture, Economic Research Service, 1999. Agriculture Information Bulletin No. 749.
- 98 National Restaurant Association (NRA). "Industry at a Glance." Accessed at <http://www.restaurant.org/research/ind_glance.cfm> on April 12, 2002.
- 99 Lin B, Guthrie J, Frazao E. *Away-From-Home Foods Increasingly Important to Quality of American Diet*. Washington, DC: U.S. Department of Agriculture, Economic Research Service, 1999. Agriculture Information Bulletin No. 749.
- 100 Zoumas-Morse C, Rock CL, Sobo EJ, Neuhouser ML. "Children's Patterns of Macronutrient Intake and Associations with Restaurant and Home Eating." *Journal of the American Dietetic Association* 2001, vol. 101, pp. 923-925.
- 101 Pereira, MA, et al. "Fast-Food Habits, Weight Gain, and Insulin Resistance (The CARDIA Study): 15-year Prospective Analysis." *Lancet* 2005, vol. 365, pp. 36-42.
- 102 Thompson OM, et al. "Food Purchased Away from Home as a Predictor of Change in BMI z-score among Girls." *International Journal of Obesity* 2004, vol. 28, pp. 282-289.
- 103 Binkley JK, et al. "The Relation between Dietary Change and Rising U.S. Obesity." *International Journal of Obesity* 2000, vol. 24, pp. 1032-1039.
- 104 Jeffery RW, French SA. "Epidemic Obesity in the United States: Are Fast Food and Television Viewing Contributing?" *American*

Journal of Public Health 1998, vol. 88, pp. 277-280.

105 McCrory MA, Fuss PJ, Saltzman E, Roberts SB. "Dietary Determinants of Energy Intake and Weight Regulation in Healthy Adults." *Journal of Nutrition* 2000, vol. 130 (Supplement), pp. 276S-279S.

106 McCrory MA, Fuss PJ, Hays NP, Vinken AG, Greenberg AS, Roberts SB. "Overeating in America: Associations between Restaurant Food Consumption and Body Fatness in Healthy Adult Men and Women Ages 19 to 80." *Obesity Research* 1999, vol. 7, pp. 564-571.

107 Zoumas-Morse C, Rock CL, Sobo EJ, Neuhauser ML. "Children's Patterns of Macronutrient Intake and Associations with Restaurant and Home Eating." *Journal of the American Dietetic Association* 2001, vol. 101, pp. 923-925.

108 Clemens LH, et al. "The Effect of Eating Out on Quality of Diet in Premenopausal Women." *Journal of the American Dietetic Association* 1999, vol. 99, pp. 422-444.

109 Schmidt M, et al. "Fast-Food Intake and Diet Quality in Black and White Girls." *Archives of Pediatric and Adolescent Medicine* 2004, vol. 159, pp. 626-631

110 S.A. Bowman and B.T. Vinyard. "Fast-Food Consumers vs. Non-Fast-Food Consumers: A Comparison of Their Energy Intakes, Diet Quality, and Overweight Status." *Journal of the American College of Nutrition* 2004, vol. 23, pp. 163-168.

111 S. Paeratakul, et al. "Fast-Food Consumption among U.S. Adults and Children: Dietary and Nutrient Intake Profile." *Journal of the American Dietetic Association* 2003, vol. 103, pp. 1332-1338.

112 Jeffery RW, French SA. "Epidemic Obesity in the United States: Are Fast Food and Television Viewing Contributing?" *American Journal of Public Health* 1998, vol. 88, pp. 277-280.

113 French SA, Story M, Neumark-Sztainer D, Fulkerson JA, Hannan P. "Fast Food Restaurant Use among Adolescents: Associations with Nutrient Intake, Food Choices and Behavioral and Psychosocial Variables." *International Journal of Obesity* 2001, vol. 25, pp. 1823-1833.

114 McNutt SW, Hu Y, Schreiber GB, Crawford PB, Obarzanek E, Mellin L. "A Longitudinal Study of the Dietary Practices of Black and White Girls 9 and 10 Years Old at Enrollment: The NHLBI Growth and Health Study." *Journal of Adolescent Health* 1997, vol. 20, pp. 27-37.

115 Lin B, Guthrie J, Frazao E. *Away-From-Home Foods Increasingly Important to Quality of American Diet*. Washington, DC: U.S. Department of Agriculture, Economic Research Service, 1999. Agriculture Information Bulletin No. 749.

116 Jeffery RW, French SA. "Epidemic Obesity in the United States: Are Fast Food and Television Viewing Contributing?" *American Journal of Public Health* 1998, vol. 88, pp. 277-280.

117 Ma Y, Bertone ER, Stanek III EJ, Reed GW, Hebert JR, Cohen NL, Merriam PA, Ockene IS. "Association between Eating Patterns and Obesity in a Free-living US Adult Population." *American Journal of Epidemiology* 2003, vol. 158, pp. 85-92.

118 McCrory MA, Fuss PJ, Hays NP, Vinken AG, Greenberg AS, Roberts SB. "Overeating in America: Associations between Restaurant Food Consumption and Body Fatness in Healthy Adult Men and Women Ages 19 to 80." *Obesity Research* 1999, vol. 7, pp. 564-571.

119 Jacobson MF, Hurley JG. *Restaurant Confidential*. New York, NY: Workman Publishing, 2002.

120 Burton S, Creyer EH, Kees J, Huggins K. "Attacking the Obesity Epidemic: An Examination of the Potential Health Benefits of

Nutrition Information Provision in Restaurants.” American Journal of Public Health, 2006, forthcoming.

121 Johnson WG, Corrigan SA, Schlundt DG, Dubbert PM. “Dietary Restraint and Eating Behavior in the Natural Environment.” Addictive Behaviors 1990, vol. 15, pp. 285-290.

122 California Center for Public Health Advocacy. Statewide poll on March 20-31, 2007 conducted by Field Research Corporation of 523 registered California voters. Accessed at www.publichealthadvocacy.org/menulabelingpoll.html on June 20, 2007.

123 End Hunger Connecticut. State-wide poll conducted between April 17 and April 23, 2007 by the Center for Survey Research and Analysis at the University of Connecticut of 501 Connecticut residents. Accessed at www.endhungerct.org/PDF/pollresults.pdf on June 20, 2007.

124 Backstrand J, Wootan MG, Young LR, Hurley J. Fat Chance. Washington, DC: Center for Science in the Public Interest, 1997.

125 Jacobson MF, Hurley JG. Restaurant Confidential. New York, NY: Workman Publishing, 2002.

126 Schab DW, Trinh N-H T. Do artificial food colorings promote hyperactivity in children with hyperactive syndromes? A meta-analysis of double-blind placebo-controlled trials. J Dev Behav Pediatr. 2004;25:423-34.

127 Bateman B, Warner JO, Hutchinson E, et al. The effects of a double blind, placebo controlled, artificial food colourings and benzoate preservative challenge on hyperactivity in a general population sample of preschool children. Archives of Disease in Childhood. 2004;89:506-11. McCann D, Barrett A, Cooper A et al. Food additives and hyperactive behaviour in 3-year-old and 8/9-year-old children in the community: a randomized, double-blinded, placebo-controlled trial. Lancet. 2007(Nov 3);370:1560-7.

128 See www.cspinet.org/new/200806022.html.

129 UK Food Standards Agency. www.food.gov.uk/news/newsarchive/2007/sep/foodcolours.

130 Poulter S. Food firms junk danger additives. Daily Mail. Sept. 8, 2007.
www.dailymail.co.uk/pages/live/articles/news/news.html?in_article_id=480644&in_page_id=1770.

131 www.foodstandards.gov.uk/news/newsarchive/2007/sep/additivesboard.

132 www.europarl.europa.eu/sides/getDoc.do?pubRef=-//EP//TEXT+TA+P6-TA-2008-0330+0+DOC+XML+V0//EN.

133 www.food.gov.uk/multimedia/pdfs/coloursletter.pdf.

134 Some dyes also have been shown to cause allergic somatic reactions.

135 R. Hingson and D. Kenkel, "Social and Health Consequences of Underage Drinking," in Reducing Underage Drinking: A Collective Responsibility, Background Papers (Washington, DC: The National Academies Press, 2004).

136 Grant, BF. Estimates of US children exposed to alcohol abuse and dependence in the family. American Journal of Public Health, Vol 90, Issue 1, pp.112-115.

137 Brown, S.A., and Tapert, S. F. Health consequences of adolescent alcohol involvement. In: National Research Council and Institute of Medicine, Bonnie, R.J., and O'Connell, M.E., eds. Reducing Underage Drinking: A Collective Responsibility. Washington, DC: National Academies Press, 2004. pp. 383-401.

138 Brown, Sandra A. et al. "A Developmental Perspective on Alcohol and Youths 16 to 20 Years of Age." Pediatrics Vol. 121 Supplement April 2008, pp. S290-S310.

139 Medina KL, Schweinsburg AD, Cohen-Zion M, Nagel BJ, & Tapert SF (2007). "Effects of alcohol and combined marijuana and alcohol use during adolescence on hippocampal volume and asymmetry." *Neurotoxicology & Teratology*, 29, 141-152.

140 Zeigler, Donald W. et al. "The neurocognitive effects of alcohol on adolescents and college students." *Preventive Medicine*, Volume 40, Issue 1, January 2005, Pages 23-32.

141 Table 2. Projections of the Population by Selected Age Groups and Sex for the United States: 2010 to 2050 (NP2008-T2). Source: Population Division, U.S. Census Bureau. Release Date: August 14, 2008.

142 NIAAA Alcohol Alert, No. 40, April 1998.

143 NIAAA Alcohol Alert, No. 74, January 2008.

144 See cited research in: Brody, Jane E. "Query for Aging Patients: How Much Do You Drink?" *New York Times*, Section D pg.7. December 16, 2008.

145 10th Special Report to Congress, pages 3, 240.